

# Heavy periods (menstrual bleeding)

This information sheet has been given to you to help answer some of the questions you may have about heavy periods and the treatments that are available. This leaflet provides information on heavy periods only. If you have other symptoms, such as bleeding in-between periods or bleeding after sex, please mention these in your consultation. If you have any further questions, please speak to a doctor or nurse caring for you.

## What are heavy periods?

A woman may be described as having 'heavy periods' if she has excessive menstrual bleeding over several menstrual cycles in a row that interfere with her physical, emotional and social quality of life. The amount of blood that is lost during a woman's period varies considerably from one individual to another, making it difficult to give a general description of heavy periods. The amount of blood lost can also vary at different times in a woman's life and in relation to operations and medications.

## What are the signs and symptoms of heavy periods?

A woman with heavy periods can experience one or more of the signs and symptoms described below:

- Blood loss with your periods that requires you to change your sanitary wear (pads or tampons) frequently (every one to two hours or more) especially when you are using large sanitary pads, super plus tampons or night time towels.
- Having to wear two pads or a pad and a tampon at the same time to control the flow.
- Periods that last more than seven days.
- Passing blood clots during periods. The size of clots can vary from the size of a 5p to the size of a 2p coin, or larger.
- Feeling tired and drained during and just after your period.
- A diagnosis of anaemia.
- 'Flooding'— when bleeding is so heavy that seating or sheets become soaked with blood while sitting or lying down.
- Feeling that your periods are interfering in your everyday life.

## What causes heavy periods?

In women 25–40 years of age, 80% with heavy periods have no underlying cause. However, the conditions listed below can sometimes be the cause of heavy periods:

- **Endometrial polyps or uterine fibroids** – Benign (non-cancerous) growths in the womb (uterus). Fibroids are usually bigger than polyps.
- **Endometriosis:** When the cells that line the inside of the uterus (the endometrium) are present elsewhere outside the uterus, such as on the ovaries or fallopian tubes. This may cause heavy periods associated with period pain, painful sex and chronic pelvic pain.

- **Intra-uterine device (IUD):** The coil inserted into the womb as a method of contraception can increase blood loss during periods by 40–50%.
- **Chronic pelvic inflammatory disease (PID):** Ongoing infection in the pelvic region of the body.
- **Polycystic ovarian syndrome (PCOS):** A condition of having multiple cysts in the ovaries, causing heavy, irregular periods, and sometimes an increase in weight and facial hair..
- **Endometrial hyperplasia:** When the endometrium (inner lining of the womb) is thicker than usual in response to an excess of oestrogen.
- **Coagulation disorders:** Blood clotting disorders, for example, von Willebrand disease.
- **Hypothyroidism:** An under-active thyroid gland.
- Some other **disease conditions**, such as liver or renal disease.
- **Medications:** There are some medications (such as warfarin) which are used to thin blood, and which can make periods heavier.
- **Cancer of the lining of the uterus:** This is very rare, but can also cause heavy periods.

## Do I need any tests to confirm the diagnosis?

There are a number of tests that can be carried out to find the cause of heavy periods. These are normally taken after you have had a full assessment with a member of your healthcare team.

### 1. Medical history

Your clinician (a doctor or a specialist nurse) will ask some questions about your concerns, medical history, the nature of your bleeding and any other related symptoms that you may have. You will be asked:

- about your menstrual cycle
  - when your last period was
  - how many days your period lasts
  - how often your periods come
  - how often you change your sanitary pads
  - what type of sanitary pads you use
  - if you have pain
  - whether or not you experience flooding
  - how your periods affect your everyday life.
- About your medication
  - if you have taken any medications for your periods and whether these have helped or not
  - if there are any other medications that you are currently taking for conditions other than your heavy periods.
- Whether you experience any other sort of vaginal bleeding such as bleeding between periods and bleeding after sexual intercourse.
- Whether you experience pelvic pain during your periods and/or pelvic pain during sexual intercourse.
- About what type of contraception you use and have used in the past. The number of pregnancies you have had and your desire to have future children.

- When you last had a cervical smear test and whether the results were normal.
- Whether you bleed for a prolonged period after sustaining a minor cut or going to the dentist.
- About your family's medical history to enable the clinician to identify whether it's possible that a hereditary condition (such as a blood clotting disorder) may be responsible for your heavy periods.

## 2. Pelvic examination

Your clinician may ask to perform a pelvic examination for a number of reasons, such as to check for an enlarged womb, which may indicate fibroids. Your clinician must obtain your consent and give you the option of having someone with you while the examination is being performed (a chaperone). A pelvic examination includes the following:

- **Vulva examination** – This involves an inspection of your external sexual organs for evidence of external bleeding and signs of infection.
- **Speculum examination of the vagina and cervix** – This uses a device called a speculum (the same instrument that is used when you go for a cervical smear test) which is gently inserted into the vagina to enable the clinician to inspect the vagina and the cervix (neck of the womb) for evidence of any abnormal changes in them.
- **Bimanual palpation** – An internal examination of your vagina, which involves the clinician inserting two fingers into the vagina while using the other hand to gently press on your abdomen. This helps to assess whether your womb or ovaries are tender or enlarged, and to identify the presence of conditions such as fibroids.

## 3. Blood test

Your clinician may take one or a number of the following blood tests:

- Full blood count – To detect iron deficiency anaemia, which is often caused by loss of iron following prolonged heavy periods. If your blood test shows that you have iron deficiency anaemia, a course of iron medicine will be prescribed for you.
- Thyroid function test (TFT) – This test may be used to detect whether or not a thyroid hormonal imbalance may be the reason for your heavy periods.
- Clotting screen – This may be done to determine whether a problem with your blood clotting mechanism may be responsible for your heavy periods.
- FSH/LH – These are hormones that control the menstrual cycles. Their levels in the blood may be checked if there has been some irregular bleeding.

The results of your blood test are usually available within one week. You will be given a date and a helpline number/email address to contact for the results of your blood test.

## 4. Pelvic scan

This is an examination of the womb using ultrasound waves to create images of structures in the pelvis. It is a painless investigation and can be done either via the abdomen (trans-abdominal scan) or via the vagina (trans-vaginal scan). The trans-vaginal scan is usually preferred as it produces better image of the uterus (womb).

## 5. Endometrial biopsy

In some cases of heavy periods where a pelvic scan shows that the endometrium (inner lining of the womb) is thicker than normal, an endometrial biopsy (a small sample of the inner lining of the womb) will be taken for laboratory tests to find out the reason for the thickness. Endometrial biopsy is obtained by inserting a straw-like tube via the vagina into the womb to obtain the sample. This procedure may cause you to have a crampy pain (like a period pain) while it is being taken and a small amount of spotting after the procedure. Endometrial biopsy can be done on its own or after a hysteroscopy.

## 6. Hysteroscopy

A hysteroscopy is a procedure which uses a fine telescope called a hysteroscope to examine the lining and shape of the uterus (womb) to look for fibroids, polyps and an overgrowth of the lining of the womb (hyperplasia). For more information please see the leaflet, **Having a hysteroscopy**.

## What treatments are available?

The treatment that is recommended for you will depend on your medical history and on the outcome of the tests and investigations that have been performed. If a reason for the bleeding (such as a fibroid) is found, then this will be treated. Sometimes, no cause for heavy periods can be found. In these cases, heavy periods can be treated by drugs or by surgery with the aim of decreasing the amount of bleeding. Some treatments may stop your periods completely and others may have an impact on your fertility. Your doctor will discuss any proposed treatments with you in detail.

The following treatments are currently recommended by the National Institute of Health and Clinical Excellence, which provides guidance on appropriate treatment for people with specific conditions:

### Drug treatments

The drug treatments below are listed in the order they will be recommended. If the first treatment is unsuitable for you, we will try the next treatment on the list. All the drug treatments can be used for as long as you need them.

1. **Levonorgestrel intrauterine system (Mirena IUS)** – This is a small plastic device that is inserted into the uterus (womb) and slowly releases a small amount of the progesterone hormone. The device is implanted during an outpatient appointment at the hospital. The Mirena IUS can reduce menstrual bleeding by up to 90%, and acts as a contraceptive as well. However, it can cause irregular bleeding when first inserted. See 'The Mirena IUS' manufacturer's leaflet for more details.
2. **Tranexamic acid tablets** – These pills can reduce the heaviness of period bleeding by 40–50% in some cases. They work by reducing the breakdown of blood clots in the uterus. Tablets are taken for three to five days during each period. Side effects are usually minor and may include stomach upset. These tablets are not a contraceptive and can be taken by women who wish to conceive. We will provide a prescription for the first four weeks and your GP will prescribe the tablets after this time.

3. **Non-steroidal anti-inflammatory drugs (NSAIDs)** – These come in various types, with Mefenamic Acid being the most commonly prescribed. NSAIDs can reduce period bleeding by 20–50%. They work by reducing the body's production of a hormone like substance, called prostaglandin, which is linked to heavy periods. NSAIDs are also painkillers and can help reduce period pain. Common side effects include indigestion and diarrhoea. These tablets are not a contraceptive and can be taken by women who wish to conceive. We will provide a prescription for the first four weeks and your GP will prescribe the tablets after this time.
4. **Combined oral contraceptive pill (COC)** – This reduces period bleeding by about 30% in some women. We will provide a prescription for the first four weeks and your GP will prescribe the tablets after this time.
5. **Long-acting progesterone contraceptives** – This includes the contraceptive injection and the contraceptive implant. They are helpful in treating heavy periods, as most women do not have any period bleeding after a few months. However, some women can have prolonged bleeding. We will supply and administer the first dose, and your GP will then continue to prescribe this treatment moving forward. Please see the Family Planning Association's leaflets, entitled, Your guide to contraceptive injections and Your guide to the contraceptive implant.
6. **Oral progesterone** – This is taken two to three times a day from days five to 26 of your menstrual cycle, counting the first day of your period as day one. It works by preventing the endometrium (inner lining of the womb) from growing quickly. Some common, short-term side effects of oral progesterone include weight gain, bloating, breast tenderness, headaches. Oral progesterone is not licensed as a contraceptive, but may have some effect on preventing conception. This means that it may not be a suitable option for women looking to conceive. It also means that additional methods of contraception should be used to protect against pregnancy. We will provide a prescription for the first four weeks; your GP will prescribe the tablets after this time.
7. **Gonadorelin analogues** – These medications work by putting you into a temporary induced menopause, and therefore stopping the menstrual cycle. They are not suitable for long-term use, as there are menopausal side effects and they can also lead to thinning of the bones (osteoporosis). We will supply and administer your first prescription, and your GP will prescribe this treatment moving forward.

## **Surgical treatments**

The clinician will explain what surgical treatment will entail. Leaflets explaining each of the different surgeries are available to help you make your choice.

Surgical treatments are only considered when medical treatments are not effective, and should only be undertaken when there is no desire for a future pregnancy. Below is a list of the surgical options available.

1. **Endometrial ablation** – This involves the destruction of the endometrium (inner lining of the womb). To do this, an instrument is passed through the vagina into the uterine cavity. There are different techniques available. The aim is to prevent the endometrium from thickening.

2. **Hysterectomy** – The surgical removal of the uterus (womb). Hysterectomy is not commonly used to treat heavy periods. For more details, please see our leaflet, **Having an abdominal hysterectomy - an operation to remove your womb**.
3. If you have fibroids there are **other surgical treatments**, such as fibroid removal, myomectomy or uterine artery embolisation. Please see our leaflet, **Surgery to remove your polyps or fibroids** for more information.

## What happens if I do not get treatment?

Not treating heavy periods is an option if there is no serious reason for your heavy bleeding and your periods do not interfere too much with your normal life. However, if the cause of your heavy period is not treated, continued heavy periods over a length of time could lead to you having a very low blood iron level (anaemia) which can lead to dizziness, breathlessness, tiredness and occasionally sudden collapse that may require a blood transfusion.

## Is there anything I can do to help?

Whether or not you decide to have surgical or medical treatment, having a balanced diet that is low in fat and contains food items that have a high iron content, such as red meat, spinach and fortified breakfast cereal, will help ensure your blood iron level is within normal limits. Other healthy lifestyle habits, such as having regular exercise and avoiding smoking, high intake of alcohol and using recreational drugs will keep you healthy.

## Contact us

If you have any questions or concerns about heavy periods, please contact the McNair Centre on **020 7188 3585** (Monday to Thursday from 9am to 5pm, and Fridays from 9am to 4.45pm).

For more information leaflets on conditions, procedures, treatments and services offered at our hospitals, please visit [www.guysandstthomas.nhs.uk/leaflets](http://www.guysandstthomas.nhs.uk/leaflets)

**Women's Health Concern** – An independent service to advise, reassure and educate women about their health concerns. Located at Whitehall House, 41 Whitehall, London SW1A 2BY.

**t:** 0845 123 2319 (local rate) **w:** [www.womens-health-concern.org](http://www.womens-health-concern.org)

**Patient Advice and Liaison Service (PALS)** – To make comments or raise concerns about the Trust's services, please contact PALS. Ask a member of staff to direct you to the PALS office or:  
**t:** 020 7188 8801 at St Thomas' **t:** 020 7188 8803 at Guy's **e:** [pals@gstt.nhs.uk](mailto:pals@gstt.nhs.uk)

**Language support services** – If you need an interpreter or information about your care in a different language or format, please get in touch using the following contact details.  
**t:** 020 7188 8815 **fax:** 020 7188 5953

**NHS Choices** – Provides online information and guidance on all aspects of health and healthcare, to help you make choices about your health.  
**w:** [www.nhs.uk](http://www.nhs.uk)

**Leaflet number: 2462/VER3**

Date published: August 2014

Review date: August 2017

© 2014 Guy's and St Thomas' NHS Foundation Trust