

# HLA incompatible kidney transplantation

This leaflet explains more about HLA (Human Leukocyte Antigen) incompatible transplantation which is sometimes referred to as tissue type incompatible transplantation. This leaflet includes information on the benefits, risks and alternatives, and what you can expect when you come to hospital.

**If you have any further questions, please speak to a doctor or nurse caring for you.**

You will have already been given information about the different types of kidney transplants. The aim of this leaflet is to explain more about HLA incompatible kidney transplantation, which is different from 'routine' transplantation in a number of ways. The transplant team here at Guy's and St Thomas' have performed over 50 transplants of this kind, making us one of the most experienced teams in the UK for this sort of transplantation. We are conducting important research in this area, so you may be asked to participate in a research study. Participation is entirely voluntary, and deciding that you do not want to be involved will not affect your clinical care in any way.

There are two forms of incompatibility:

- Blood group incompatibility is when the donor and recipient are different blood groups.
- HLA tissue type incompatibility is when there are antibodies against the donor's tissue type.

This leaflet explains more about HLA incompatible kidney transplantation.

For more information about blood group incompatible transplants, please ask for a copy of our leaflet: **"Blood group incompatible kidney transplantation"**

## What is HLA incompatible transplantation?

Sometimes a patient waiting to have a kidney transplant has antibodies in their blood that would react against the donor kidney and damage it. The antibodies in the transplant recipient are said to be 'incompatible' with the donor kidney. If your tissue type is incompatible with your donor, this means that you have previously been exposed to cells or tissue from another person and that you have formed antibodies that can attack tissue from your donor (in this case, the donated kidney). Most people are exposed as a result of a previous blood transfusion, pregnancy (for women) or previous organ transplantation.

Routine transplantation in these circumstances would result in severe rejection and potential loss of the kidney. However, it is possible to remove or suppress these antibodies in a process known as 'desensitisation', which can allow the transplant to take place.

In order to increase the chances of a successful outcome for HLA incompatible transplants, specific treatments (plasma exchange or immunoadsorption) alongside medication are available to combat any antibodies you may have formed against your donor.

## **What are the risks of HLA incompatible transplantation?**

There are some risks that are associated with transplant surgery, regardless of whether you are compatible or incompatible with your donor. These risks include the risk of bleeding (both during and after the operation), the risk of wound infection and the risk of death. There is more detail about the risks associated with transplant surgery in chapter 2 of ["Your guide to kidney transplantation"](#). Please ask for a copy of this booklet if you don't have one.

There are additional risks associated with an HLA incompatible transplant, which include:

### **Rejection**

There is an increased risk of rejection after HLA incompatible transplantation. For compatible transplants, around 25% will experience some rejection over the first year. For HLA incompatible transplants this can be up to 40%.

There are two types of rejection: antibody-mediated rejection, and cell-mediated rejection.

With HLA incompatible transplants, the most common and risky type of rejection is antibody-mediated rejection. The risks are highest within the first month after the transplant. If you have good kidney function for the first month following surgery, the risk of rejection decreases.

An early antibody-mediated rejection can be difficult to treat, and may require more plasma exchange and additional medication. If the rejection does not respond to treatment, the transplant could fail.

Cell-mediated rejection tends to be easier to treat with intravenous steroids.

### **Bleeding**

All surgery carries the risk of bleeding, but when we remove antibodies with plasma exchange, we also remove some of the factors in your blood that are important for making your blood clot. This could increase your risk of bleeding during and after transplantation. It may be necessary to replace these factors before the surgery.

### **Infections**

All transplant patients are at increased risk of infections due to the immunosuppressive medications that we give. For patients who have an HLA incompatible transplant, we tend to prescribe stronger immunosuppressants, and this may increase your risk of infection. Sometimes these infections can be very serious, and may even be fatal. You will be closely monitored after the transplant for signs of infection.

### **Patient survival**

Although all types of transplant surgery carry the risk of death, this risk is increased with HLA transplantation. This is partly due to the greater risk of infection associated with taking stronger immunosuppressive medications, which are necessary to help prevent rejection of the donor kidney.

If you have been dependent on dialysis for a long time, you may have other health problems which reduce your ability to survive a major complication, such as a serious infection.

## Survival of the transplant

If you are compatible with your donor, there is a 10% risk that your transplanted kidney will stop working within the first five years after transplantation. If you are HLA incompatible, the risk that your donor kidney stops working within the first five years is 30–40%.

## Are there any alternatives?

All of this may sound worrying, but the chances are that if you are at the stage of considering an HLA incompatible transplant, you have exhausted all possible compatible donors. You may already have been entered into the **National Kidney Donor Sharing Scheme** where you and your donor are paired with a donor and recipient in another centre to improve the chances of a better matched kidney for both recipients. You have the option to remain in the National Kidney Donor Sharing Scheme and/or on the national waiting list for a deceased donor transplant. All patients are different and we recommend you discuss your options with your doctor if you have any questions or concerns.

HLA incompatible transplantation is risky, but it is often considered better in the long term to be transplanted, even with a high-risk transplant, than to remain on dialysis.

## Giving my consent (permission)

We want to involve you in decisions about your care and treatment. If you decide to go ahead, you will be asked to sign a consent form. This states that you agree to have the treatment and you understand what it involves.

## Antibody removal: How can I prepare for the treatment?

In order for the treatment to take place, we need to have access to your blood – this is known as vascular access. If you are receiving regular haemodialysis treatment, you will probably already have vascular access, which can be used. If you are not receiving haemodialysis, you will need to have a dialysis line inserted. Most people will need more than one antibody removal treatment and will need a tunnelled haemodialysis line to be inserted. If you only need one treatment then a temporary haemodialysis line can be used. For more information about these procedures, please ask us for a copy of our leaflets, ["Having a tunnelled haemodialysis line inserted"](#), or ["Having a non-tunnelled haemodialysis line inserted"](#)

## What happens during the treatment?

The treatment will take place on Astley Cooper Dialysis Unit, 5<sup>th</sup> Floor, Borough Wing, Guy's Hospital, and will be carried out by specialist nurses. Plasma exchange usually takes two to three hours, so you may wish to bring a book, ipod, laptop or tablet with you to help pass the time. You may also wish to bring a snack. If you are receiving regular dialysis, we will dialyse you after the plasma exchange.

You will receive treatment using a machine that is very similar to a dialysis machine. The machine will filter your blood and remove the plasma (the liquid component of blood), which contains your antibodies. The plasma will then be replaced with albumin (a type of protein) or a similar product – we call this **plasma exchange**. The number of treatments required depends on the levels of antibodies in your blood.

Some patients will undergo a technique called **immunoadsorption** instead of plasma exchange. In this case, the blood is passed through a column that binds antibodies. The blood and plasma are then returned to the patient.

These treatments will be given during the days leading up to your transplant, and the type of treatment (plasma exchange or immunoadsorption) will depend on the amount of antibodies in your blood. In some cases, with low antibodies, no treatment will be necessary, despite the antibody incompatibility.

## After the treatment

Occasionally people feel tired or light-headed after having plasma exchange or immunoadsorption. You should eat breakfast or lunch on the day of treatment as this can help to prevent or minimise this feeling. You will also need to arrange for a relative or friend to accompany you home.

We will carry out a final cross-match against your donor the day before your transplant and your antibodies will need to be checked on the morning of the surgery. If we haven't been able to reduce the antibody level sufficiently, it may be necessary to postpone the transplant at short notice. If this is the case, we usually carry out further treatments to try to reduce the antibodies and hopefully the surgery can take place at a later date.

## Immunosuppressive medicines

You will be given anti-rejection medication both before and after your kidney transplant – these are called immunosuppressants, because they work by inhibiting or preventing the activity of the immune system. All patients who undergo kidney transplantation will need to take anti-rejection medication for the rest of the time that the kidney is working. You can read more about these medications in our booklet, ["Your guide to kidney transplantation"](#). If you have not already been given a copy, please ask us for one.

Patients who are incompatible with their donor usually require higher dose of anti-rejection medication than those undergoing a routine transplant. These usually include **tacrolimus** and **mycophenolate**, which you will be asked to start taking two to seven days before your transplant, depending on your degree of incompatibility. This is to ensure that there is an adequate level of the anti-rejection drug in your blood at the time of surgery.

You will also be given some additional medicines prior to the transplant that will further help to reduce the risk of rejection. These may include:

- Alemtuzumab
- Antithymocyte immunoglobulin

**Alemtuzumab** and **antithymocyte immunoglobulin** inhibit the cells of your immune system that are involved in rejection.

- Alemtuzumab is given by subcutaneous (under the skin) injection during your surgery and a second dose may be given on the first day following surgery.
- Antithymocyte immunoglobulin is given by intravenous infusion on the day of your transplant and then daily for three days after.

The use of alemtuzumab in transplantation is "unlicensed". This means that, although it has been used in transplantation for many years, the manufacturer's licence for the product is for a different condition or range of conditions. More information on the use of unlicensed medicines can be found in our leaflet, **Unlicensed medicines: A guide for patients**. If you have not already been given a copy, please ask us for one.

We hope you have found this leaflet useful. All patients are different and we strongly advise that you discuss your situation carefully with your medical and nursing teams to ensure that you understand the implications for you personally.

## Contact us

Please contact us if you have further queries:

<b>Mr Nizam Mamode</b> (consultant transplant surgeon)	<b>020 7188 1543</b>
<b>Lisa Silas</b> (advanced nurse practitioner – living donation)	<b>020 7188 5688</b>
<b>Irmen Generalao</b> (clinical nurse specialist – antibody removal)	<b>020 7188 9276</b>

**Out of hours, please contact 07917084532.**

For more information leaflets on conditions, procedures, treatments and services offered at our hospitals, please visit [www.guysandstthomas.nhs.uk/leaflets](http://www.guysandstthomas.nhs.uk/leaflets)

## Pharmacy Medicines Helpline

If you have any questions or concerns about your medicines, please speak to the staff caring for you or call our helpline.

**t:** 020 7188 8748 9am to 5pm, Monday to Friday

## Your comments and concerns

For advice, support or to raise a concern, contact our Patient Advice and Liaison Service (PALS). To make a complaint, contact the complaints department.

<b>t:</b> 020 7188 8801 (PALS)	<b>e:</b> <a href="mailto:pals@gstt.nhs.uk">pals@gstt.nhs.uk</a>
<b>t:</b> 020 7188 3514 (complaints)	<b>e:</b> <a href="mailto:complaints2@gstt.nhs.uk">complaints2@gstt.nhs.uk</a>

## Language Support Services

If you need an interpreter or information about your care in a different language or format, please get in touch using the following contact details.

**t:** 020 7188 8815      **fax:** 020 7188 5953

## NHS 111

Offers medical help and advice from fully trained advisers supported by experienced nurses and paramedics. Available over the phone 24 hours a day.

**t:** 111

## NHS Choices

Provides online information and guidance on all aspects of health and healthcare, to help you make choices about your health.

**w:** [www.nhs.uk](http://www.nhs.uk)

## Get involved and have your say: become a member of the Trust

Members of Guy's and St Thomas' NHS Foundation Trust contribute to the organisation on a voluntary basis. We count on them for feedback, local knowledge and support. Membership is free and it is up to you how much you get involved. To find out more, and to become a member:

**t:** 0800 731 0319    **e:** [members@gstt.nhs.uk](mailto:members@gstt.nhs.uk)    **w:** [www.guysandstthomas.nhs.uk/membership](http://www.guysandstthomas.nhs.uk/membership)

**Leaflet number: 4254/VER1**

Date published: June 2016

Review date: June 2019

© 2014 Guy's and St Thomas' NHS Foundation Trust