

Vardenafil

Male Cardiovascular Health Clinic

This information sheet aims to answer some of the questions you may have about using vardenafil to treat your erectile dysfunction. Please ensure that you read the manufacturer's leaflet that accompanies your supply of vardenafil for further details on the treatment. Vardenafil is not available for use on the NHS in South East London and will require a private prescription. If you have any further questions or concerns, please speak to a doctor or nurse caring for you.

What is vardenafil?

Vardenafil is an effective treatment for most men who suffer from erectile dysfunction (ED or impotence). It works by allowing the normal process of an erection to occur.

Vardenafil does not change your sex drive and is only recommended for men with erectile dysfunction. Vardenafil will not work without sexual stimulation of some kind.

How do I use vardenafil?

Vardenafil can be taken with or without food, although if you take it with a heavy meal it could delay how quickly it works.

You should take vardenafil about one hour before anticipated sexual activity. It is effective from 30 minutes to approximately two hours after taking it.

Do not take more than one tablet in any 24 hour period. Do not exceed the dose that your doctor has prescribed for you. Do not drink grapefruit juice when you take vardenafil, as this can affect how well the drug works.

What are the side effects of using vardenafil?

Possible side effects include:

- headache
- flushing
- dizziness
- indigestion
- nasal congestion
- temporary changes in your vision, such as colour disturbance, blurred vision or increased perception of light. Please bear this in mind if you are planning to drive. Please contact your doctor immediately and stop taking vardenafil if you experience a sudden decrease or loss of vision.

The side effects may reduce over time but please talk to your doctor or nurse if you have any questions about them. Further information on side effects can be found in the manufacturer's patient information leaflet that comes with the medicine.

Things to avoid

Vardenafil must not be taken in combination with **long-acting** nitrates, such as isosorbide mononitrate (sometimes known as Imdur[®], Ismo[®], Elantan[®]) or nicorandil, as this can cause a sudden and dangerous drop in blood pressure.

Is there anything else I need to know?

If you are taking **short-acting** nitrates, either as glyceryl trinitrate (GTN) spray or as tablets under the tongue, you should only use vardenafil after you have talked it through carefully with your doctor or nurse.

Do not take a dose of vardenafil within 12 hours of a dose of GTN. This means that if you need to use your GTN spray you will have to wait until 12 hours after you have taken vardenafil. If you have chest pain that does not go away with rest, go to hospital and tell them that you have taken vardenafil.

Never take vardenafil with non-prescription drugs, such as amyl nitrate/nitrates (poppers). If you are unsure whether you take nitrates please consult with your doctor or nurse.

Vardenafil may also interact with other medicines, so let your nurse know immediately if there are any changes to the medicines that you take (including those that you buy yourself or herbal/homeopathic medicines) so that they can make sure that this treatment is still appropriate.

You can continue to eat foods that contain nitrates, such as green leafy vegetables and root vegetables.

Chest pain following the use of vardenafil

Before prescribing vardenafil, your doctor will have made a thorough assessment to ensure that it is safe for you to take this drug. **However, in the unlikely event that you experience chest pain after using vardenafil:**

- you must rest
- seek medical advice if the pain doesn't go away within 10–15 minutes.

If you need to seek medical help, you must let the doctors know that you have taken vardenafil, as this will affect the treatment you receive.

You must not use any form of nitrate therapy – including the spray or tablets under the tongue – without medical advice.

Where can I get a repeat prescription?

You can get a repeat prescription from your GP.

Useful sources of information

Contact us

If you or your partner has any questions about vardenafil, please do not hesitate to call the cardiac nurse specialist, **t:** 07900 224890. Please note that the specialist nurse may not be able to return your call immediately but will aim to do so as soon as possible.

For more information leaflets on conditions, procedures, treatments and services offered at our hospitals, please visit www.guysandstthomas.nhs.uk/leaflets

Pharmacy Medicines Helpline

If you have any questions or concerns about your medicines, please speak to the staff caring for you or call our helpline.

t: 020 7188 8748 9am to 5pm, Monday to Friday

Your comments and concerns

For advice, support or to raise a concern, contact our Patient Advice and Liaison Service (PALS). To make a complaint, contact the complaints department.

t: 020 7188 8801 (PALS) **e:** pals@gstt.nhs.uk

t: 020 7188 3514 (complaints) **e:** complaints2@gstt.nhs.uk

Language and accessible support services

If you need an interpreter or information about your care in a different language or format, please get in touch.

t: 020 7188 8815 **e:** languagesupport@gstt.nhs.uk

NHS 111

Offers medical help and advice from fully trained advisers supported by experienced nurses and paramedics. Available over the phone 24 hours a day.

t: 111

NHS Choices

Provides online information and guidance on all aspects of health and healthcare, to help you make choices about your health.

w: www.nhs.uk

Get involved and have your say: become a member of the Trust

Members of Guy's and St Thomas' NHS Foundation Trust contribute to the organisation on a voluntary basis. We count on them for feedback, local knowledge and support. Membership is free and it is up to you how much you get involved. To find out more, please get in touch.

t: 0800 731 0319 **e:** members@gstt.nhs.uk **w:** www.guysandstthomas.nhs.uk/membership

Leaflet number: 2523/VER5

Date published: September 2018

Review date: September 2021

© 2018 Guy's and St Thomas' NHS Foundation Trust

A list of sources is available on request